

*This example is set up in BrushScriptX-Italic. It uses:*

```
\usepackage[T1]{fontenc}
%\usepackage{pbsi}
\renewcommand{\rmdefault}{pbsi}
\renewcommand{\mddefault}{xl}
\renewcommand{\bfdefault}{xl}
\usepackage[defaultmathsizes,noasterisk]{mathastext}
\begin{document}\boldmath
```

*Typeset with mathastext 1.13 (2011/03/11).*

To illustrate some Hilbert Space properties of the co-Poisson summation, we will assume  $K = 2$ . The components  $(a_v)$  of an adele  $a$  are written  $a_p$  at finite places and  $a_r$  at the real place. We have an embedding of the Schwartz space of test-functions on  $\mathcal{R}$  into the Bruhat-Schwartz space on  $\mathcal{A}$  which sends  $\psi(x)$  to  $\varphi(a) = \prod_p 1_{|a_p|_p \leq 1}(a_p) \cdot \psi(a_r)$ , and we write  $\mathcal{E}'_{\mathcal{R}}(g)$  for the distribution on  $\mathcal{R}$  thus obtained from  $\mathcal{E}'(g)$  on  $\mathcal{A}$ .

**Theorem 1.** Let  $g$  be a compact Bruhat-Schwartz function on the ideles of 2. The co-Poisson summation  $\mathcal{E}'_{\mathcal{R}}(g)$  is a square-integrable function (with respect to the Lebesgue measure). The  $L^2(\mathcal{R})$  function  $\mathcal{E}'_{\mathcal{R}}(g)$  is equal to the constant  $-\int_{\mathcal{A}^\times} g(u) |u|^{-1/2} d^*u$  in a neighborhood of the origin.

*Proof.* We may first, without changing anything to  $\mathcal{E}'_{\mathcal{R}}(g)$ , replace  $g$  with its average under the action of the finite unit ideles, so that it may be assumed invariant. Any such compact invariant  $g$  is a finite linear combination of suitable multiplicative translates of functions of the type  $g(u) = \prod_p 1_{|u_p|_p = 1}(u_p) \cdot f(u_r)$  with  $f(t)$  a smooth compactly supported function on  $\mathcal{R}^\times$ , so that we may assume that  $g$  has this form. We claim that:

$$\int_{\mathcal{A}^\times} |\varphi(u)| \sum_{g \in 2^\times} |g(gu)| \sqrt{|u|} d^*u < \infty$$

Indeed  $\sum_{g \in 2^\times} |g(gu)| = |f(|u|)| + |f(-|u|)|$  is bounded above by a multiple of  $|u|$ . And  $\int_{\mathcal{A}^\times} |\varphi(u)| |u|^{3/2} d^*u < \infty$  for each Bruhat-Schwartz function on the adeles (basically, from  $\prod_p (1 - p^{-3/2})^{-1} < \infty$ ). So

$$\mathcal{E}'(g)(\varphi) = \sum_{g \in 2^\times} \int_{\mathcal{A}^\times} \varphi(u) g(gu) \sqrt{|u|} d^*u - \int_{\mathcal{A}^\times} \frac{g(u)}{\sqrt{|u|}} d^*u \int_{\mathcal{A}} \varphi(x) dx$$

$$\mathcal{E}'(g)(\varphi) = \sum_{g \in 2^\times} \int_{\mathcal{A}^\times} \varphi(u/g) g(u) \sqrt{|u|} d^*u - \int_{\mathcal{A}^\times} \frac{g(u)}{\sqrt{|u|}} d^*u \int_{\mathcal{A}} \varphi(x) dx$$

Let us now specialize to  $\varphi(a) = \prod_p 1_{|a_p|_p \leq 1}(a_p) \cdot \psi(a_r)$ . Each integral can be evaluated as an infinite product. The finite places

contribute 0 or 1 according to whether  $q \in \mathbb{Z}^\times$  satisfies  $|q|_p < 1$  or not. So only the inverse integers  $q = 1/n$ ,  $n \in \mathbb{Z}$ , contribute:

$$\mathcal{E}'_{\mathcal{R}}(q)(\psi) = \sum_{n \in \mathbb{Z}^\times} \int_{\mathcal{R}^\times} \psi(nt) f(t) \sqrt{|t|} \frac{dt}{2|t|} - \int_{\mathcal{R}^\times} \frac{f(t)}{\sqrt{|t|}} \frac{dt}{2|t|} \int_{\mathcal{R}} \psi(x) dx$$

We can now revert the steps, but this time on  $\mathcal{R}^\times$  and we get:

$$\mathcal{E}'_{\mathcal{R}}(q)(\psi) = \int_{\mathcal{R}^\times} \psi(t) \sum_{n \in \mathbb{Z}^\times} \frac{f(t/n)}{\sqrt{|n|}} \frac{dt}{2\sqrt{|t|}} - \int_{\mathcal{R}^\times} \frac{f(t)}{\sqrt{|t|}} \frac{dt}{2|t|} \int_{\mathcal{R}} \psi(x) dx$$

Let us express this in terms of  $\alpha(q) = (f(q) + f(-q))/2\sqrt{|q|}$ :

$$\mathcal{E}'_{\mathcal{R}}(q)(\psi) = \int_{\mathcal{R}} \psi(q) \sum_{n \geq 1} \frac{\alpha(q/n)}{n} dq - \int_0^\infty \frac{\alpha(q)}{q} dq \int_{\mathcal{R}} \psi(x) dx$$

So the distribution  $\mathcal{E}'_{\mathcal{R}}(q)$  is in fact the even smooth function

$$\mathcal{E}'_{\mathcal{R}}(q)(q) = \sum_{n \geq 1} \frac{\alpha(q/n)}{n} - \int_0^\infty \frac{\alpha(q)}{q} dq$$

As  $\alpha(q)$  has compact support in  $\mathcal{R} \setminus \{0\}$ , the summation over  $n \geq 1$  contains only vanishing terms for  $|q|$  small enough. So  $\mathcal{E}'_{\mathcal{R}}(q)$  is equal to the constant  $-\int_0^\infty \frac{\alpha(q)}{q} dq = -\int_{\mathcal{R}^\times} \frac{f(q)}{\sqrt{|q|}} \frac{dq}{2|q|} = -\int_{\mathcal{A}^\times} g(t)/\sqrt{|t|} d^*t$  in a neighborhood of 0. To prove that it is  $\mathcal{L}^2$ , let  $\beta(q)$  be the smooth compactly supported function  $\alpha(1/q)/2|q|$  of  $q \in \mathcal{R}$  ( $\beta(0) = 0$ ). Then ( $q \neq 0$ ):

$$\mathcal{E}'_{\mathcal{R}}(q)(q) = \sum_{n \in \mathbb{Z}} \frac{1}{|q|} \beta\left(\frac{n}{q}\right) - \int_{\mathcal{R}} \beta(q) dq$$

From the usual Poisson summation formula, this is also:

$$\sum_{n \in \mathbb{Z}} \gamma(nq) - \int_{\mathcal{R}} \beta(q) dq = \sum_{n \neq 0} \gamma(nq)$$

where  $\gamma(q) = \int_{\mathcal{R}} \exp(i2\pi q\omega) \beta(\omega) d\omega$  is a Schwartz rapidly decreasing function. From this formula we deduce easily that  $\mathcal{E}'_{\mathcal{R}}(q)(q)$  is itself in the Schwartz class of rapidly decreasing functions, and in particular it is square-integrable.  $\square$

*It is useful to recapitulate some of the results arising in this proof:*

*Theorem 2. Let  $g$  be a compact Bruhat-Schwartz function on the ideles of  $\mathbb{Z}$ . The co-Poisson summation  $\mathcal{E}'_{\mathcal{R}}(g)$  is an even function on  $\mathbb{R}$  in the Schwartz class of rapidly decreasing functions. It is constant, as well as its Fourier Transform, in a neighborhood of the origin. It may be written as*

$$\mathcal{E}'_{\mathcal{R}}(g)(y) = \sum_{n \geq 1} \frac{\alpha(y/n)}{n} - \int_0^{\infty} \frac{\alpha(y)}{y} dy$$

*with a function  $\alpha(y)$  smooth with compact support away from the origin, and conversely each such formula corresponds to the co-Poisson summation  $\mathcal{E}'_{\mathcal{R}}(g)$  of a compact Bruhat-Schwartz function on the ideles of  $\mathbb{Z}$ . The Fourier transform  $\int_{\mathbb{R}} \mathcal{E}'_{\mathcal{R}}(g)(y) \exp(i2\pi\omega y) dy$  corresponds in the formula above to the replacement  $\alpha(y) \mapsto \alpha(1/y)/|y|$ .*

*Everything has been obtained previously.*