

This example is set up in Helvetica with Fourier. It uses:

```
\usepackage[T1]{fontenc}
\usepackage[scaled]{helvet}
\usepackage{fourier}
\renewcommand{\rmdefault}{phv}
\usepackage[italic,defaultmathsizes,noasterisk]{mathastext}
```

Typeset with mathastext 1.13 (2011/03/11).

To illustrate some Hilbert Space properties of the co-Poisson summation, we will assume $K = \mathbf{Q}$. The components (a_v) of an adèle a are written a_p at finite places and a_r at the real place. We have an embedding of the Schwartz space of test-functions on \mathbf{R} into the Bruhat-Schwartz space on \mathbf{A} which sends $\psi(x)$ to $\varphi(a) = \prod_p \mathbf{1}_{|a_p|_p \leq 1}(a_p) \cdot \psi(a_r)$, and we write $E_{\mathbf{R}}'(g)$ for the distribution on \mathbf{R} thus obtained from $E'(g)$ on \mathbf{A} .

Theorem 1. *Let g be a compact Bruhat-Schwartz function on the ideles of \mathbf{Q} . The co-Poisson summation $E_{\mathbf{R}}'(g)$ is a square-integrable function (with respect to the Lebesgue measure). The $L^2(\mathbf{R})$ function $E_{\mathbf{R}}'(g)$ is equal to the constant $-\int_{\mathbf{A}^\times} g(v)|v|^{-1/2} d^*v$ in a neighborhood of the origin.*

Proof. We may first, without changing anything to $E_{\mathbf{R}}'(g)$, replace g with its average under the action of the finite unit ideles, so that it may be assumed invariant. Any such compact invariant g is a finite linear combination of suitable multiplicative translates of functions of the type $g(v) = \prod_p \mathbf{1}_{|v_p|_p=1}(v_p) \cdot f(v_r)$ with $f(t)$ a smooth compactly supported function on \mathbf{R}^\times , so that we may assume that g has this form. We claim that:

$$\int_{\mathbf{A}^\times} |\varphi(v)| \sum_{q \in \mathbf{Q}^\times} |g(qv)| \sqrt{|v|} d^*v < \infty$$

Indeed $\sum_{q \in \mathbf{Q}^\times} |g(qv)| = |f(|v|)| + |f(-|v|)|$ is bounded above by a multiple of $|v|$. And $\int_{\mathbf{A}^\times} |\varphi(v)| |v|^{3/2} d^*v < \infty$ for each Bruhat-Schwartz function on the adeles (basically, from $\prod_p (1 - p^{-3/2})^{-1} < \infty$). So

$$\begin{aligned} E'(g)(\varphi) &= \sum_{q \in \mathbf{Q}^\times} \int_{\mathbf{A}^\times} \varphi(v) g(qv) \sqrt{|v|} d^*v - \int_{\mathbf{A}^\times} \frac{g(v)}{\sqrt{|v|}} d^*v \int_{\mathbf{A}} \varphi(x) dx \\ E'(g)(\varphi) &= \sum_{q \in \mathbf{Q}^\times} \int_{\mathbf{A}^\times} \varphi(v/q) g(v) \sqrt{|v|} d^*v - \int_{\mathbf{A}^\times} \frac{g(v)}{\sqrt{|v|}} d^*v \int_{\mathbf{A}} \varphi(x) dx \end{aligned}$$

Let us now specialize to $\varphi(a) = \prod_p \mathbf{1}_{|a_p|_p \leq 1}(a_p) \cdot \psi(a_r)$. Each integral can be evaluated as an infinite product. The finite places contribute 0 or 1 according to whether $q \in \mathbf{Q}^\times$ satisfies $|q|_p < 1$ or not. So only the inverse integers $q = 1/n$, $n \in \mathbf{Z}$, contribute:

$$E_{\mathbf{R}}'(g)(\psi) = \sum_{n \in \mathbf{Z}^\times} \int_{\mathbf{R}^\times} \psi(nt) f(t) \sqrt{|t|} \frac{dt}{2|t|} - \int_{\mathbf{R}^\times} \frac{f(t)}{\sqrt{|t|}} \frac{dt}{2|t|} \int_{\mathbf{R}} \psi(x) dx$$

We can now revert the steps, but this time on \mathbf{R}^\times and we get:

$$E_{\mathbf{R}}'(g)(\psi) = \int_{\mathbf{R}^\times} \psi(t) \sum_{n \in \mathbf{Z}^\times} \frac{f(t/n)}{\sqrt{|n|}} \frac{dt}{2\sqrt{|t|}} - \int_{\mathbf{R}^\times} \frac{f(t)}{\sqrt{|t|}} \frac{dt}{2|t|} \int_{\mathbf{R}} \psi(x) dx$$

Let us express this in terms of $\alpha(y) = (f(y) + f(-y))/2\sqrt{|y|}$:

$$E'_{\mathbf{R}}(g)(\psi) = \int_{\mathbf{R}} \psi(y) \sum_{n \geq 1} \frac{\alpha(y/n)}{n} dy - \int_0^{\infty} \frac{\alpha(y)}{y} dy \int_{\mathbf{R}} \psi(x) dx$$

So the distribution $E'_{\mathbf{R}}(g)$ is in fact the even smooth function

$$E'_{\mathbf{R}}(g)(y) = \sum_{n \geq 1} \frac{\alpha(y/n)}{n} - \int_0^{\infty} \frac{\alpha(y)}{y} dy$$

As $\alpha(y)$ has compact support in $\mathbf{R} \setminus \{0\}$, the summation over $n \geq 1$ contains only vanishing terms for $|y|$ small enough. So $E'_{\mathbf{R}}(g)$ is equal to the constant $-\int_0^{\infty} \frac{\alpha(y)}{y} dy = -\int_{\mathbf{R}^{\times}} \frac{f(y)}{\sqrt{|y|}} \frac{dy}{2|y|} = -\int_{\mathbf{A}^{\times}} g(t)/\sqrt{|t|} d^*t$ in a neighborhood of 0. To prove that it is L^2 , let $\beta(y)$ be the smooth compactly supported function $\alpha(1/y)/2|y|$ of $y \in \mathbf{R}$ ($\beta(0) = 0$). Then ($y \neq 0$):

$$E'_{\mathbf{R}}(g)(y) = \sum_{n \in \mathbf{Z}} \frac{1}{|y|} \beta\left(\frac{n}{y}\right) - \int_{\mathbf{R}} \beta(y) dy$$

From the usual Poisson summation formula, this is also:

$$\sum_{n \in \mathbf{Z}} \gamma(ny) - \int_{\mathbf{R}} \beta(y) dy = \sum_{n \neq 0} \gamma(ny)$$

where $\gamma(y) = \int_{\mathbf{R}} \exp(i2\pi yw) \beta(w) dw$ is a Schwartz rapidly decreasing function. From this formula we deduce easily that $E'_{\mathbf{R}}(g)(y)$ is itself in the Schwartz class of rapidly decreasing functions, and in particular it is square-integrable. \square

It is useful to recapitulate some of the results arising in this proof:

Theorem 2. *Let g be a compact Bruhat-Schwartz function on the ideles of \mathbf{Q} . The co-Poisson summation $E'_{\mathbf{R}}(g)$ is an even function on \mathbf{R} in the Schwartz class of rapidly decreasing functions. It is constant, as well as its Fourier Transform, in a neighborhood of the origin. It may be written as*

$$E'_{\mathbf{R}}(g)(y) = \sum_{n \geq 1} \frac{\alpha(y/n)}{n} - \int_0^{\infty} \frac{\alpha(y)}{y} dy$$

with a function $\alpha(y)$ smooth with compact support away from the origin, and conversely each such formula corresponds to the co-Poisson summation $E'_{\mathbf{R}}(g)$ of a compact Bruhat-Schwartz function on the ideles of \mathbf{Q} . The Fourier transform $\int_{\mathbf{R}} E'_{\mathbf{R}}(g)(y) \exp(i2\pi wy) dy$ corresponds in the formula above to the replacement $\alpha(y) \mapsto \alpha(1/y)/|y|$.

Everything has been obtained previously.